

2. Sectional Committee on Nuclear Accident, Committee on Comprehensive Synthetic Engineering, Science Council of Japan (2014), A review of the model comparison of transportation and deposition of radioactive materials released to the environment as a result of the Tokyo Electric Power Company's Fukushima Daiichi Nuclear Power Plant accident, available at www.jpgeu.org/scj/report/20140902scj_report_e.pdf, accessed during May 2017.
3. Ohara S., A.Mukai, N.Fujiwara (2015), The deposition map obtained through radioactivity analysis of mosses, *Kankyō Gijyū*, 44, pp.460-467 (Japanese version).
4. Sanada Y., T.Sugita, Y.Nishizawa, A.Kondo, T.Torii (2014), The aerial radiation monitoring in Japan after the Fukushima Daiichi nuclear power plant accident, *Progress in Nuclear Science and Technology*, 4, pp.76-80.
5. Saito K., I.Tanihata, M.Fujiwara, T.Saito, S.Shimoura, T.Otsuka, Y.Onda, M.Hoshi, Y.Ikeuchi, F.Takahashi, N.Kinouchi, J.Saegusa, A.Seki, H.Takemiya, T.Shibata (2015), Detailed deposition density maps constructed by large-scale soil sampling for gamma-ray emitting radioactive nuclides from the Fukushima Daiichi Nuclear Power Plant accident, *Journal of Environmental Radioactivity*, 139, pp.308-319.
6. Koizumi A., T.Niisoe, K.H.Harada, Y.Fujii, A.Adachi, T.Hitomi, H.Ishikawa (2013), ¹³⁷Cs trapped by biomass within 20km of the Fukushima Daiichi Nuclear Power Plant, *Environmental Science and Technology*, 47, pp.9612-9618.
7. Steinnes E., O.Njastad (1993), Use of mosses and lichens for regional mapping of ¹³⁷Cs fallout from the Chernobyl accident, *Journal of Environmental Radioactivity*, 21, pp.65-73.
8. Kato H., Y.Onda, M.Teramage (2012), Depth distribution of ¹³⁷Cs, ¹³⁴Cs, and ¹³¹I in soil profile after Fukushima Dai-ichi Nuclear Power Plant Accident, *Journal of Environmental Radioactivity*, 111, pp.59-64.
9. Rigol A., M.Vidal, G.Rauret (2002), An overview of the effect of organic matter on soil-radiocaesium interaction: implications in root uptake, *Journal of Environmental Radioactivity*, 58, pp.191-216.
10. Ii I., K.Tanoi, Y.Uno, T.Nobori, A.Hirose, N.Kobayashi, N.Nihei, T.Ogawa, Y.Tao, M.Kanno, J.Nishiwaki, M.Mizoguchi (2015), Radioactive cesium concentration of lowland rice growth in the decontaminated paddy fields in Iitate-Village in Fukushima, *RADIOISOTOPES*, 64, pp.299-310 (Japanese version).
11. Nakao A., S.Funakawa, T.Kosaki (2009), Hydroxy-Al polymers block the frayed edge sites of illitic minerals in acid soils: Studies in southwestern Japan at various weathering stages, *European Journal of Soil Science*, 60, pp.127-138.
12. Kawada Y and T.Yamada (2012), Radioactivity ratio of ¹³⁴Cs/¹³⁷Cs released by the nuclear accidents, *Isotope News*, No.697, pp.16-20.